Data table for Figure 14. Uncontrolled high blood pressure among adults aged 20 and over with hypertension, by sex and age: United States, 1988–1994 through 2011–2014

Excel and PowerPoint: http://www.cdc.gov/nchs/hus/contents2016.htm#fig14

Sex and age	Uncontrolled high blood pressure among adults aged 20 and over with hypertension 1										
	1988–1994		1999–2002		2003–2006		2007–2010		2011–2014		
	Percent	SE	Percent	SE	Percent	SE	Percent	SE	Percent	SE	
	Percent of population										
20 years and over, age-adjusted ²											
Both sexes	77.2	1.4	70.6	1.7	63.3	1.9	55.8	1.8	52.8	1.9	
Men	83.2	1.6	73.3	1.8	65.0	2.3	61.4	2.1	58.1	1.9	
Women	68.5	2.2	61.8	3.4	53.6	4.1	46.3	2.7	45.5	3.1	
20 years and over, crude											
Both sexes	73.9	1.0	67.3	1.5	58.6	1.4	49.3	1.2	47.0	1.8	
Men	79.3	1.3	67.1	1.9	58.4	2.1	52.3	1.6	49.9	2.1	
Women	68.8	1.6	67.4	1.7	58.8	1.3	46.4	1.3	44.3	1.9	

2011-2014

Percent	SE		
74.1	4.6		
64.2	5.1		
49.7	3.1		
38.2	3.1		
46.5	4.4		
55.5	8.0		
37.7	3.7		
36.8	2.7		
44.5	2.8		
61.5	3.0		
	74.1 64.2 49.7 38.2 46.5 55.5 37.7 36.8 44.5		

SE is standard error.

SOURCE: NCHS, National Health and Nutrition Examination Survey. See Appendix I, National Health and Nutrition Examination Survey (NHANES).

¹Uncontrolled high blood pressure among adults with hypertension is defined as measured systolic pressure of at least 140 mm Hg or diastolic pressure of at least 90 mm Hg, among those with measured high blood pressure or reporting taking antihypertensive medication.

² Estimates are age-adjusted to the year 2000 standard population using five age groups: 20–34 years, 35–44 years, 45–54 years, 55–64 years, and 65 years and over. Age-adjusted estimates in this table may differ from other age-adjusted estimates based on the same data and presented elsewhere if different age groups are used in the adjustment procedure. See Appendix II, Age adjustment; Blood pressure, high. Also see Table 54.